



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.


Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

r the radius of the circle. Then $(x-b+r)^2 + (y-r)^2 = r^2$ is the equation to the circle. Hence the invariants are $\Delta = -4a^2$, $\Delta' = -r^2$, $\theta = -4a(b-r+a)$, $\theta' = -4a(b-r)$. In order that the circle and parabola may touch $(\theta\theta' - 9\Delta\Delta')^2 = 4(\theta^2 - 3\Delta\theta') \times (\theta'^2 - 3\Delta'\theta)$.



$\therefore 16r^5 + (13a - 48b)r^4 + 8(a^2 + 6b^2 - 14ab)r^3 + 8(15ab^2 - 2b^3 - 9a^2b)r^2 + 32ab(3ab - a^2 - 2b^2)r + 16ab^2(b^2 + a^2 - 2ab) = 0$. This gives the value of r .

Area of semi-parabola $= \frac{2}{3}b\sqrt{ab}$, area of circle $= \pi r^2$, area of portion left $= \frac{2}{3}b\sqrt{ab} - \pi r^2$.

Let A be the vertex of the parabola, B the centroid required, C the centroid of the semi-parabola, D the centroid of the circle. Let (α, β) be the coordinates of B . The coordinates of C are $(\frac{2}{3}b, \frac{2}{3}\sqrt{ab})$; of D , $(b-r, r)$.

$$\frac{CD}{BC} = \frac{HK}{BH} = \frac{GF}{EF} = \frac{\frac{2}{3}b\sqrt{ab} - \pi r^2}{\pi r^2} \text{ or } \frac{EG}{EF} = \frac{4b\sqrt{ab}}{3\pi r^2}.$$

$$\therefore \frac{b-r-a}{\frac{2}{3}b-a} = \frac{4b\sqrt{ab}}{3\pi r^2}. \therefore a = \frac{3[4b^2\sqrt{ab} + 5\pi r^3 - 5\pi br^2]}{5[4b\sqrt{ab} - 3\pi r^2]}.$$

$$\text{And } \frac{DK}{CH} = \frac{EG}{EF} = \frac{4b\sqrt{ab}}{3\pi r^2} = \frac{r-\beta}{\frac{2}{3}\sqrt{ab} - \beta}; \therefore \beta = \frac{3(ab^2 - \pi r^3)}{4b\sqrt{ab} - 3\pi r^2}.$$

208. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, Eng.

Hanging at rest over a smooth pulley are two equal scale pans of the same mass. Two equal particles, the one inelastic and the other elastic, are simultaneously dropped from the same height one into each scale pan. Show that each impact after the first must occur when the pans have returned to the *status quo ante*, and find the total space described by either pan before motion ceases.

Remark by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

This is the same as problem 121, Mechanics. A solution of this problem is found in Vol. VIII, No. 10, pp. 203-4, October, 1901.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

141. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Given that the highest factor of a prime p contained in $m!$ is p^{m-s} ; find general expressions involving p and m and s , from which, when a solution is possible, m can be determined when s is a given integer and p is a given prime. Is it then possible in any case to have more solutions than one?